

# Ophthalmic Artery Doppler Indices Changes in Preeclampsia

Mahsa Naemi<sup>1</sup>, Maasoumeh Saleh<sup>1\*</sup> , Mahboubeh Saleh<sup>2</sup>

1. Department of Obstetrics and Gynecology, Shariati Hospital, Tehran University of Medical Sciences, Tehran, Iran
2. Department of Obstetrics and Gynecology, Fasa University of Medical Sciences, Fars, Iran



## Article Info

 [10.30699/jogcr.8.2.125](https://doi.org/10.30699/jogcr.8.2.125)

**Received:** 2022/11/03;

**Accepted:** 2022/12/18;

**Published Online:** 22 Feb 2023;

Use your device to scan and read the article online



## Corresponding Information:

**Maasoumeh Saleh,**

Department of Obstetrics and Gynecology, Shariati Hospital, Tehran University of Medical Sciences, Tehran, Iran

**Email:** [Salehmaasoumeh@yahoo.com](mailto:Salehmaasoumeh@yahoo.com)

## ABSTRACT

**Background & Objective:** Preeclampsia (PE) is a major complication of pregnancy worldwide. Doppler ultrasound of maternal arteries, including uterine and ophthalmic arteries (OA), plays an important role in determining individuals progressing PE. The aim of this study is to compare OA indices in pregnant women with a diagnosis of PE and women without it.

**Materials & Methods:** This study evaluated pregnant women who had delivery in Shariati Hospitals, Tehran, Iran between January 2021 and March 2022. The maternal OA Doppler waves were obtained in preeclamptic and non-preeclamptic women at 35 to 37 weeks of gestation by a maternal-fetal medicine specialist. OA Doppler parameters including first and second PSV, second to first PSV ratio (Peak ratio, PR) and PI were obtained for each eye and also average between the eyes.

**Results:** Out of 148 included participants, 48 cases progressed to PE at 35 to 37 weeks of gestation and 100 cases did not have PE at the same gestational age. All OA parameters including (First and second peak systolic velocities (PSV), second to first PSV Ratio (PR) and Pulsatility Index (PI) in each eye; as well as, the average of these indices in both eyes) were assessed. PR and PI between the two eyes were statistically different in those with PE.

**Conclusion:** OA Doppler indices change in pregnant women with PE. These changes could help to earlier and better diagnosis of PE and prevention of maternal and fetal sequelae.

**Keywords:** Preeclampsia, Ophthalmic Artery, Doppler Indices, Pregnancy Complication



Copyright © 2023, This is an original open-access article distributed under the terms of the Creative Commons Attribution-noncommercial 4.0 International License which permits copy and redistribution of the material just in noncommercial usages with proper citation.

## Introduction

Preeclampsia (PE) is a major complication of pregnancy with adverse effects on mother and fetus. It is one of the important causes of maternal mortality. It is defined as follows, acute onset of maternal hypertension and proteinuria after 20th weeks of gestational age (1). There are several methods for predicting PE, including the following: maternal characteristics, uterine arteries (UtA) and ophthalmic arteries (OA) Doppler indices, and serum biomarkers. Impaired blood flow in maternal arteries may be observed in pregnant women before the onset of PE and at the time of diagnosis and it could be detected by Doppler ultrasound (2). Doppler study is a safe and accessible method and can be done in the least time and also in a portable way. In this study, we have assessed the changes of OA Doppler indices in pregnant women with PE.

## Methods

This study evaluated pregnant women who had delivery in Shariati Hospitals, Tehran, Iran between January 2021 and March 2022. The maternal OA Doppler waves were obtained in preeclamptic and non-preeclamptic women at 35 to 37 weeks of gestation by a maternal-fetal medicine specialist. All cases gave written informed consent to participate in this study and it was approved by the Research Ethics Committee at Tehran University of Medical Sciences.

The inclusion criteria were singleton pregnancies examined at 35 to 37 weeks of gestation. We excluded pregnant women with type 1 and 2 diabetes mellitus (DMI, II), chronic hypertension (HTN), autoimmune diseases, and those with fetal aneuploidy or abnormality. 48 cases progressed to PE at 35 to 37 weeks of gestation, and 100 cases did not have PE at the same gestational age were included in the study.

PE defined as follows: high blood pressure measurements (systolic blood pressure [SBP]  $\geq 140$ mmHg and/or diastolic pressure [DBP]  $\geq 90$

mmHg) that was associated with acute-onset proteinuria ( $\geq 1+$  protein on a dipstick analysis or  $\geq 300$  mg in urine 24 hours). Severe PE was defined as PE with severe hypertension (SBP  $\geq 160$  mmHg and/or DBP  $\geq 110$  mmHg) or signs or symptoms of end-organ damage.

Ophthalmic artery (OA) waveforms were obtained with a 7-15 MHz linear transducer that was placed

horizontally on the upper eyeball. Pulsed-wave Doppler was recorded medial to the optic nerve and three to five waveforms were obtained. OA Doppler parameters including first and second PSV, second to first PSV ratio (Peak ratio, PR) and PI were obtained for each eye and also average between the eyes (Figure 1). Finally, all analyses were performed in SPSS (version 26) and P-values less than 0.05 were defined as statistically significant.




Figure 1. A sample of ophthalmic artery Doppler waveforms

### Results

A total of 148 pregnant women were included in the study. The population contained 100 normal cases and 48 pregnancies with PE (42 cases of mild PE and 6 cases of severe PE). This study showed that the following variables were higher in preeclamptic pregnancies compared to those without PE: Right (Rt.) and left (Lt.) second PSV, average (Av.) of second PSV between the eyes, Rt. PR, Lt. PR, and Av. of PR

between the eyes (Table 1). The area under curve (AUC) values (95% CI) of three OA Doppler indices, including Rt. PR (0.96, 0.94 – 0.98), Lt. PR (0.96, 0.94 – 0.99) and Av. of PR between the eyes (0.97, 0.94 – 0.99) were higher than 0.90 (Table 2). Lt. PR, Rt. PR, Av. of PR and Av. of PI between the eyes had the most important changes in PE.

Table 1. Ophthalmic artery Doppler indices of study population of 148 pregnancies, according to development of mild and severe preeclampsia (PE)

Characteristics	Normal (n=100)	PE (n=48)			P*	P#	p†
		Total (n=48)	Mild PE (n=42)	Severe PE (n=6)			
Right First PSV	31.4 (25.5, 38.4)	24.5 (20.2, 27.4)	24.2 (20.5, 27.4)	27.0 (16.2, 31.6)	<0.001	<0.001	0.838
Right Second PSV	17.3 (13.6, 20.6)	21.3 (18.5, 26.0)	21.3 (18.5, 23.8)	25.0 (14.6, 29.3)	<0.001	<0.001	0.451
Right ophthalmic artery PI	2.0 (1.8, 2.3)	1.2 (1.0, 1.6)	1.2 (1.0, 1.6)	1.3 (1.0, 1.5)	<0.001	<0.001	0.949

Characteristics	Normal (n=100)	PE (n=48)			P*	P#	p†
		Total (n=48)	Mild PE (n=42)	Severe PE (n=6)			
Left First PSV	32.0 (25.6, 38.5)	24.4 (20.3, 27.5)	24.0 (20.5, 27.5)	27.2 (16.1, 31.6)	<0.001	<0.001	0.816
Left Second PSV	17.4 (14.0, 21.0)	21.3 (18.4, 26.4)	21.2 (18.4, 24.4)	25.1 (14.7, 29.5)	<0.001	<0.001	0.442
Left ophthalmic artery PI	2.0 (1.8, 2.3)	1.2 (1.0, 1.6)	1.2 (1.0, 1.6)	1.3 (1.1, 1.6)	<0.001	<0.001	0.997
Average PR between two eyes	53.7 (48.4, 60.3)	84.6 (76.0, 96.2)	84.0 (76.0, 94.6)	91.8 (79.8, 98.5)	<0.001	<0.001	0.220
Average of first PSV between two eyes	31.7 (25.6, 38.4)	24.4 (20.2, 27.5)	24.1 (20.5, 27.5)	27.1 (16.2, 31.6)	<0.001	<0.001	0.827
Average of second PSV between two eyes	17.4 (13.8, 20.6)	21.3 (18.5, 26.2)	21.2 (18.5, 24.1)	25.1 (14.7, 29.4)	<0.001	<0.001	0.446
Average of ophthalmic artery PI between two eyes	2.0 (1.8, 2.3)	1.2 (1.0, 1.6)	1.2 (1.0, 1.6)	1.3 (1.0, 1.6)	<0.001	<0.001	0.976
Right PR	53.8 (48.3, 60.5)	84.5 (74.7, 96.2)	84.0 (74.7, 94.6)	91.1 (79.0, 98.5)	<0.001	<0.001	0.257
Left PR	53.8 (48.3, 60.3)	84.6 (77.3, 96.1)	84.1 (77.0, 94.5)	92.4 (80.5, 98.6)	<0.001	<0.001	0.189

**Table 2.** The diagnostic measures of ophthalmic artery Doppler indices in PE

Variables	AUC (95% CI)	Best Cut point
Right First PSV	0.75 (0.64, 0.86)	0.06
Right Second PSV	0.70 (0.56, 0.84)	0.06
Right ophthalmic artery PI	0.87 (0.78, 0.97)	0.06
Left First PSV	0.75 (0.64, 0.86)	0.06
Left Second PSV	0.70 (0.57, 0.84)	0.06
Left ophthalmic artery PI	0.88 (0.79, 0.97)	0.06
Average PR between two eyes	0.97 (0.94, 0.99)	0.07
Average of first PSV between two eyes	0.75 (0.64, 0.86)	0.06
Average of second PSV between two eyes	0.70 (0.57, 0.84)	0.06
Average of ophthalmic artery PI between two eyes	0.88 (0.78, 0.97)	0.07
Right PR	0.96 (0.94, 0.98)	0.07
Left PR	0.96 (0.94, 0.99)	0.07


## Discussion

PE is responsible for more than 16% of maternal mortalities worldwide (3). Screening and early diagnosis of PE are very important issues in obstetrics medicine, especially in high-risk pregnancies (1, 4). Fetal and maternal complications of PE are serious (5).

Therefore, it is necessary to study markers for early prediction and diagnosis of PE. Recently, researches have evaluated cerebrovascular changes in preeclamptic women and some of them have evaluated OA Doppler indices in prediction and diagnosis of PE.


It is important to know more about maternal arteries Doppler changes before progressing to PE or at the time of PE.

The ophthalmic artery (OA) is the first branch of the internal carotid artery and has embryological, anatomical and functional similarities with the intracranial vasculature. Its branches supply the orbit, meninges, face, and upper nose (6). This artery is easily accessible for ultrasound and can have useful



information from the cerebral circulation that is not easily accessible for evaluation. Therefore, in some studies, ultrasonographic examination of OAs has been used to predict neurological consequences. Although, computed tomography (CT) and magnetic resonance imaging (MRI) are more difficult methods because of less access, high cost, and risks to mothers with high blood pressure during transportation, but OA Doppler can be performed in the shortest possible time and on a portable basis. Figure 2 shows OA Doppler analysis.

**Figure 2.** Shows OA Doppler indices (7). The first PSV wave is peak velocity S1, it represents the maximum velocity during systole and is as an ejection parameter in cardiac systole and augmentation of S2 is related to reflection of pressure wave. The peak diastolic velocity, (D), is the maximum velocity which rises due to vascular elastic recoil during cardiac diastole, and (d), the end diastolic velocity, represents the minimum velocity.



**Figure 3.** Shows pulse wave (PW) reflection and transmission between OA and UtA. Reflection will increase with vasoconstriction and vascular tone (8). Therefore, the reason for the increase in second PSV in OA Doppler assessment in PE is well shown in this figure. Also, the relationship between OA Doppler indices changes and PE may be related to hemodynamic adaptation to the pregnancy (9).

Most studies have shown that PR in the OA Doppler indices was the best predictor of PE and had the most important changes in PE (10, 11). Our study showed that Av. of PR between the two eyes had the significant change in PE. In some studies, both OA PI and PSV were reported higher in PE while others have shown a significant elevation in normal pregnancies (12-20). Unlike PR, first and second PSV changes of the OA are unreliable indices in prediction and diagnosis of PE. Therefore, it suggests against first and second PSV indices use as routine markers for prediction and early diagnosis of PE (11, 21-23).

The strengths of this study were a higher sample size compared to other studies. OA Doppler ultrasound is especially helpful in prediction and also early diagnosis of PE. Facilitating earlier diagnosis of PE can help prevent serious maternal and fetal complications, especially maternal neurological sequelae (24).

## Conclusion

This study assessed the OA Doppler indices changes in pregnant women with PE and showed that these changes could help to earlier and better diagnosis of PE and prevention of maternal and fetal sequelae.

## Acknowledgments

None.

## Conflict of Interest

The authors declare no known conflicts of interest.

## References

1. Roberts JM, August PA, Bakris G, Barton JR, Bernstein IM, Druzin M, et al. Hypertension in pregnancy: executive summary. *Obstet Gynecol.* 2013;122(5):1122-31. [PMID] [DOI:10.1097/01.AOG.0000437382.03963.88]
2. J Spaan J, A Brown M. Renin-angiotensin system in pre-eclampsia: everything old is new again. *Obstet Med.* 2012;5(4):147-53. [PMID] [PMCID] [DOI:10.1258/om.2012.120007]
3. Khan KS, Wojdyla D, Say L, Gülmezoglu AM, Van Look PF. WHO analysis of causes of maternal death: a systematic review. *The lancet.* 2006;367(9516):1066-74. [PMID] [DOI:10.1016/S0140-6736(06)68397-9]
4. Lu JF, Nightingale CH. Magnesium sulfate in eclampsia and pre-eclampsia: pharmacokinetic principles. *Clin Pharmacokinet.* 2000;38(4):305-14. [DOI:10.2165/00003088-200038040-00002] [PMID]
5. Shahraki Z, Rezaie Keikhaie K, Amjadi N, Hashemi Bonjar Z, Jahantigh H, Doosti F, et al. Correlation of 4 Hour Urine Samples with 24-Hour Urine Samples for the Diagnosis of Preeclampsia. *J Obstet Gynecol Cancer Res.* 2022;2(2). [DOI:10.5812/jogcr.11730]
6. Bird B, Stawicki SP. *Anatomy, Head and Neck, Ophthalmic Arteries.* StatPearls Publishing, Treasure Island (FL); 2022.
7. Rasyada AR, Azhim A. Flow velocity in common carotid artery. *Carotid Artery Gender and Health* 2019.
8. Gonser M. Hemodynamic relationship between ophthalmic artery and uterine artery in pre-eclampsia: pulse wave reflection and transmission might provide the missing link. *Ultrasound Obstet Gynecol.* 2019;53(1):135-6. [DOI:10.1002/uog.20121] [PMID]
9. Kalafat E, Laoreti A, Khalil A, Da Silva Costa F, Thilaganathan B. Ophthalmic artery Doppler for prediction of pre-eclampsia: systematic review and meta-analysis. *Ultrasound Obstet Gynecol.* 2018;51(6):731-7. [DOI:10.1002/uog.19002] [PMID]
10. Sarno M, Wright A, Vieira N, Sapantzoglou I, Charakida M, Nicolaidis KH. Ophthalmic artery Doppler in prediction of pre-eclampsia at 35-37 weeks' gestation. *Ultrasound Obstet Gynecol.* 2020;56(5):717-24. [DOI:10.1002/uog.22184] [PMID]
11. Matias DS, Santos R, Ferreira T, Matias BS, Correia LCL. Predictive value of ophthalmic artery Doppler velocimetry in relation to hypertensive disorders of pregnancy. *J Clin Ultrasound.* 2020;48(7):388-95. [DOI:10.1002/jcu.22823] [PMID]
12. Sapantzoglou I, Wright A, Arozena MG, Campos RV, Charakida M, Nicolaidis KH. Ophthalmic artery Doppler in combination with other biomarkers in prediction of pre-eclampsia at 19-23 weeks' gestation. *Ultrasound Obstet Gynecol.* 2021;57(1):75-83. [DOI:10.1002/uog.23528] [PMID]
13. Ozdemir ME, Demirci O, Ozturkmen HA, Ulusoy NB, Ohanoglu K, Cilingir IU. What Is the Role of the Maternal Ophthalmic and Cervical Internal Carotid Arteries in Predicting Maternal Adverse Outcomes in Preeclampsia? *J Ultrasound Med.* 2020;39(8):1527-35. [DOI:10.1002/jum.15241] [PMID]
14. Onwudiegwu C, Adekanmi A, Olusanya B, Lawal O, Adedokun B, Morhason-Bello I, et al. Case-control study on ocular changes and ophthalmic Doppler velocimetric indices among preeclamptic and normotensive pregnant women in Ibadan, Nigeria. *BMJ Open Ophthalmol.* 2020;5(1). [DOI:10.1136/bmjophth-2020-000550] [PMID] [PMCID]
15. Matias DS, Costa RF, Matias BS, Cláudio Lemos Correia L. Doppler velocimetry of the orbital vessels in pregnancies complicated by preeclampsia. *J Clin Ultrasound.* 2012;40(9):576-85. [DOI:10.1002/jcu.21949] [PMID]
16. Gurgel Alves JA, Praciano de Sousa PC, Bezerra Maia EHMS, Kane SC, da Silva Costa F. First-trimester maternal ophthalmic artery Doppler analysis for prediction of pre-eclampsia. *Ultrasound Obstet Gynecol.* 2014;44(4):411-8. [DOI:10.1002/uog.13338] [PMID]
17. Hata T, Senoh D, Hata K, Kitao M. Ophthalmic artery velocimetry in pregnant women. *Lancet.* 1992;340(8812):182-3. [DOI:10.1016/0140-6736(92)93268-R] [PMID]
18. Olatunji RB, Adekanmi AJ, Obajimi MO, Roberts OA, Ojo TO. Maternal ophthalmic artery Doppler velocimetry in pre-eclampsia in Southwestern

- Nigeria. *Int J Womens Health*. 2015;7:723-34. [DOI:10.2147/IJWH.S86314] [PMID] [PMCID]
19. Anderson UD, Olsson MG, Kristensen KH, Åkerström B, Hansson SR. Review: Biochemical markers to predict preeclampsia. *Placenta*. 2012;33:S42-S7. [DOI:10.1016/j.placenta.2011.11.021] [PMID]
  20. Albaiges G, Missfelder-Lobos H, Lees C, Parra M, Nicolaides KH. One-stage screening for pregnancy complications by color Doppler assessment of the uterine arteries at 23 weeks' gestation. The authors thank Paul Seed, Division of Primary Care and Health Sciences, Guy's, King's and St. Thomas's Hospitals, for valuable statistical input. *Obstet Gynecol*. 2000;96(4):559-64. [DOI:10.1016/S0029-7844(00)00946-7] [DOI:10.1097/00006250-200010000-00015] [PMID]
  21. Andrietti S, Carlucci S, Wright A, Wright D, Nicolaides KH. Repeat measurements of uterine artery pulsatility index, mean arterial pressure and serum placental growth factor at 12, 22 and 32 weeks in prediction of pre-eclampsia. *Ultrasound Obstet Gynecol*. 2017;50(2):221-7. [DOI:10.1002/uog.17403] [PMID]
  22. Steel SA, Pearce JM, McParland P, Chamberlain GVP. Early doppler ultrasound screening in prediction of hypertensive disorders of pregnancy. *The Lancet*. 1990;335(8705):1548-51. [DOI:10.1016/0140-6736(90)91376-L] [PMID]
  23. Sigel B. A Brief History of Doppler Ultrasound in the Diagnosis of Peripheral Vascular Disease. *Ultrasound Med Biol*. 1998;24(2):169-76. [DOI:10.1016/S0301-5629(97)00264-0] [PMID]
  24. English FA, Kenny LC, McCarthy FP. Risk factors and effective management of preeclampsia. *Integr Blood Press Control*. 2015;8:7-12. [DOI:10.2147/IBPC.S50641] [PMID] [PMCID]

#### How to Cite This Article:

Naemi, M., Saleh, M., Saleh, M. Ophthalmic Artery Doppler Indices Changes in Preeclampsia Dear Colleague. *J Obstet Gynecol Cancer Res*. 2023; 8(2):125-30.

#### Download citation:

[BibTeX](#) | [RIS](#) | [EndNote](#) | [Medlars](#) | [ProCite](#) | [Reference Manager](#) | [RefWorks](#)