

# Unusual Feature of Mature Cystic Teratoma: Multiple Intra-cystic Spherical Structures: A Case report

Sedigheh Ghasemian Dizaj Mehr<sup>1\*</sup> , Robabeh Bahadori<sup>2</sup>

1. Department of Obstetrics and Gynecology, School of Medicine, Urmia University of Medical Science. Urmia, Iran
2. Reproductive Health Research Center, Clinical Research Institute, Urmia University of Medical Sciences, Urmia, Iran



## Article Info

 [10.30699/jogcr.8.6.637](https://doi.org/10.30699/jogcr.8.6.637)

**Received:** 2022/09/10;

**Accepted:** 2023/01/09;

**Published Online:** 11 Nov 2023;

Use your device to scan and read the article online



## Corresponding Information:

**Sedigheh Ghasemian Dizaj Mehr,**  
Department of Obstetrics and Gynecology,  
School of Medicine, Urmia University of  
Medical Science. Urmia, Iran

**Email:** [ghasemian.s@umsu.ac.ir](mailto:ghasemian.s@umsu.ac.ir)



Copyright © 2023, This is an original open-access article distributed under the terms of the Creative Commons Attribution-noncommercial 4.0 International License which permits copy and redistribution of the material just in noncommercial usages with proper citation.

## ABSTRACT

Ovarian mature cystic teratomas (OMCTs) due to their heterogeneous histologic entity, have a wide spectrum of radiologic features. A type of floating spherical globule that has been called “Sack of marble” or “Coins-in-sack,” but without a fat or calcification component that results in misdiagnosis on computed tomography (CT), is a rare finding. The patient was a 35-year-old healthy woman with abdominal pain and sonographic findings based on OMCT with the Coins-in-sack sign, whereas the CT manifestation was a unilocular huge cyst suggesting an adnexal cystadenoma. She had a final diagnosis of the Coins-in-sack variant of OMCT but with keratin balls, not common fat balls. Understanding the unconventional radiologic and histologic manifestations of OMCTs, is necessary to make an accurate diagnosis.

**Keywords:** Ovarian Mature Cystic Teratoma (OMCT), Floating balls, Coins-in-sack, Computed Tomography (CT)


## Introduction

Ovarian mature cystic teratoma (OMCT) is the most common benign ovarian germ cell neoplasm. It is a kind of predominantly cystic tumor that is composed of mature tissue of ectodermal, mesodermal and endodermal derivatives that result in characteristic gross appearance and histologic gross appearance and histologic with various radiologic features due to heterogeneous entity (1, 2).

Although different imaging manifestations of OMCTs, are well - described in the literature, the less common variants of this mass should be considered more, until the rare, specific radiologic features of OMCTs, are not misdiagnosed with the neoplasm. We describe a case of OMCT with an unusual imaging feature of multiple intracystic floating spherical structures.

## Case Presentation


A 35-year-old healthy woman with gravida 3 para 3 was referred to our oncology clinic due to recent abdominal pain and mass. On pelvic examination, there is a huge mobile mass in the midline of the pelvis that is extended up to the umbilical part. Pelvic transvaginal ultrasound revealed a huge 15×20 cm in mean diameter, complex cystic mass with a thick wall, including numerous mobile spherical echogenic intracystic structures without vascularization on color Doppler, that made a suggestion of a mature pelvic teratoma with the radiologic feature of the “Coins-in-sack or “Sack of marble” sign (Figure 1).



**Figure 1.** Pelvic ultrasound shows, floating intra cystic echogenic balls with flat-fluid level and scanty echogenic linear structures.

On computed tomography (CT) without contrast, it is reported large 15×20 cm unilocular cystic mass in midline of pelvic, pushing to surrounding tissues


without calcification and fat density, suggested adnexal cystadenoma ([Figure 2](#)).



**Figure 2.** CT without contrast: huge unilocular cystic mass without calcification and fat density.

Serum tumor markers were, CA-125:20 u/ml and CA-19-9: 2 u/ml. The patient underwent laparotomy. Intraoperatively right salpingo- oophorectomy was carried out due to 15×20 cm unilocular cystic mass

without adhesion or spillage. A surgical specimen cut-section showed numerous 2-10 mm brown pasty globules floating in a claylike fluid with a small amount of hair ([Figure 3](#)).



**Figure 3.** Surgical specimen cut-section with numerous pasty globules and small amount of hair.

The patient had an uneventful postoperative recovery. Permanent pathology was reported, 15×15cm mature cystic teratoma with 0.8cm wall thickness, numerous colored solid globules measuring 0.2 to 1cm in diameter, floating in turbid light brown colored fluid made of keratin material and a raised protuberance in the cyst wall (Rokitansky nodule) composed of hair tuft measuring 3 cm in the greatest diameter.

## Discussion

Ovarian mature cystic teratomas (OMCTs), because of their various compositions due to pluripotential germ cell origin, can have a wide spectrum of radiologic imaging features. In gross appearance, it is straightforward to diagnose OMCTs. Frequently, the cut section of OMCTs is unilocular cyst filled with sebaceous material, because of the wall that is lined by the squamous epithelium. It is also called a dermoid cyst, due to the presence of most skin elements, such as hair and keratin (3-5).

Fat is present in more than 93% of OMCTs and teeth or other calcification occurs in 56% of cases (2, 4). Due to the variety of appearance and histologic types in this kind of tumor, a wide spectrum of radiologic imaging features has been reported. Most OMCTs can be diagnosed by ultrasonography, but unusual types lead to misdiagnosis. Computed tomography (CT) and magnetic resonance (MR) imaging, made the diagnosis of OMCTs due to the more sensitive nature of those modalities for fat calcification (6, 7).

One of the bizarre radiologic features of OMCTs is intracystic floating balls, which have been called coins-in-sack or the "sack of marble" sign. Firstly, it had been reported on the basis of pathologic findings (8), whereas the radiologic report of this variant was done subsequently (9).

Pathologically, these floating balls were formed by the aggregation of variable sebaceous material, debris, keratin, hair, hemosiderin and fat in enough space of large cyst (6, 10, 11).

The different contents and density of these globules, made them gravity-dependent or gravity-independent in large cyst (10).

It is clear that, being essential, enough space for ball formation needs to increase with age due to the slow growth of OMCTs at an average rate of 1.8 mm each year (2, 10).

Floating globules that have been called fat balls in some studies, don't always consist of fatty components (6, 11, 12).

Sometimes, due to the lack of fat components in the floating balls, and being close to the liquid density of these spherical structures, this kind of floating balls or coins-in-sack sign, on CT modality, may be not revealed but with attention to showing different signals

for different components in MRI, it can be determined that this unusual type of OMCTs with an absence or scary amount of fat in the globules (13).

In our case, the feature of coins-in-sacks was first reported on sonography.

MCTs are usually detected easily on sonography, and their specificity is 99% (2), but the variety of OMCTs composition results in limitless sonographic appearances. Therefore, the presentation of uncommon case reports is associated with an increased accuracy of diagnosis. Despite some studies based on the large size and diameter of these floating balls, this case included small balls 0.5 to 0.8 cm in diameter. According to previous studies based on floating ball formation, a large cyst that consequently happened in older patients, whereas our patients were not much older, but the size of the cyst was huge.

These balls could be misdiagnosed with papillary projections or nodular growth, particularly in limited amounts, of malignant masses on sonography, but similar to our case, color flow mapping showed no vascularization in these balls. In our case, on CT imaging, incorrectly, it was suggested that there was a probable huge adnexal cystadenoma due to the absence of fat or calcification content.

Pathologically, these globules were made of keratin materials with no fat component, therefore, the density of this content is close to the liquid density occupying the cavity of this large cyst, so it is difficult to distinguish of these kinds of unusual features of OMCTs on CT examination. Because of these pitfalls of CT, it is preferred in questionable sonographic cases or for confirming sonographic suggestions initially.

We perform MRI examinations because we have different signals for different histologic components (13).

## Conclusion

OMCTs due to their heterogeneous histologic component, have various and challenging radiologic features, sometimes with unconventional manifestations that are not presented in the fat or calcification component, so, physicians should be aware of their distinct and uncommon appearances in different imaging modalities.

Therefore, we should elucidate uncommon findings, so that they can be helpful in training, making accurate diagnoses, and expanding our knowledge.

## Acknowledgments

The authors appreciate all the staff members of the obstetrics and gynecology department of Kowsar Hospital, Urmia, Iran for their help and suggestions.

## Informed Consent

Written informed consent was obtained from the patient.

## Ethics Committee Approval

This study was approved by the Ethics Committee of Urmia University Medical Sciences.

## Conflict of Interest

There is no conflict of interest.

## Funding and Support

There is no funding or support.

## References

- DiSaia PJ, Creasman WT, Mannell RS, McMeekin S, Mutch DG. Clinical gynecologic oncology e-book: Elsevier Health Sciences; 2017.
- Outwater EK, Siegelman ES, Hunt JL. Ovarian teratomas: tumor types and imaging characteristics. *Radiographics*. 2001;21(2):475-90. [PMID] [DOI:10.1148/radiographics.21.2.g01mr09475]
- Jagtap SV, Kshirsagar NS, Jagtap SS, Boral S, Nasre N. Ovarian teratomas: clinicopathological study at tertiary care center. *Int j Reprod Contracept Obstet Gynecol*. 2019;8(8):3318-23. [DOI:10.18203/2320-1770.ijrcog20193558]
- Sahin H, Abdullazade S, Sanci M. Mature cystic teratoma of the ovary: a cutting edge overview on imaging features. *Insights into imaging*. 2017; 8(2):227-41. [DOI:10.1007/s13244-016-0539-9] [PMID] [PMCID]
- Saba L, Guerriero S, Sulcis R, Virgilio B, Melis G, Mallarini G. Mature and immature ovarian teratomas: CT, US and MR imaging characteristics. *Eur J Radiol*. 2009;72(3):454-63. [DOI:10.1016/j.ejrad.2008.07.044] [PMID]
- Tongsong T, Wanapirak C, Khunamornpong S, Sukpan K. Numerous intracystic floating balls as a sonographic feature of benign cystic teratoma: report of 5 cases. *J Ultrasound Med*. 2006;25(12):1587-91. [DOI:10.7863/jum.2006.25.12.1587] [PMID]
- Occhipinti KA, Frankel SD, Hricak H. The ovary. *Computed tomography and magnetic resonance imaging*. *Radiol Clin North Am*. 1993;31(5):1115-32. [DOI:10.1016/S0033-8389(22)00359-1] [PMID]
- DeAngelis CE. An ovarian dermoid cyst with numerous sebum balls. *Am J Obstet Gynecol*. 1953;66(2):443-5. [PMID] [DOI:10.1016/0002-9378(53)90587-0]
- Muramatsu Y, Moriyama N, Takayasu K, Nawano S, Yamada T. CT and MR imaging of cystic ovarian teratoma with intracystic fat balls. *J Comput Assist Tomogr*. 1991;15(3):528-9.
- Şahin H, Akdoğan AI, Ayaz D, Karadeniz T, Sancı M. Utility of the "floating ball sign" in diagnosis of ovarian cystic teratoma. *Turk J Obstet Gynecol*. 2019;16(2):118. [PMID] [PMCID] [DOI:10.4274/tjod.galenos.2019.67209]
- Jantarsaengaram S, Siricharoenchai S, Vairojanavong K. Cystic ovarian teratoma with intracystic fat balls. *Ultrasound Obstet Gynecol*. 2003;22(1):102-3. [DOI:10.1002/uog.181] [PMID]
- Gol M, Saygili U, Uslu T, Erten O. Mature cystic teratoma with intracystic fat balls in a postmenopausal woman. *Eur J Obst Gynec Reprod Biol*. 2005;119(1):125-6. [DOI:10.1016/j.ejogrb.2004.06.003] [PMID]
- Wang A, Shu Y. Mature ovarian cystic teratoma with "sack of marbles" appearance on magnetic resonance imaging: A case report. *Medicine*. 2019;98(31):e1669. [PMID] [PMCID] [DOI:10.1097/MD.0000000000001669]

## How to Cite This Article:

Ghasemian Dizaj Mehr, S., Bahadori, R. Unusual Feature of Mature Cystic Teratoma: Multiple Intra-cystic Spherical Structures: A Case report. *J Obstet Gynecol Cancer Res*. 2023; 8(6):637-40.

Download citation:

[RIS](#) | [EndNote](#) | [Mendeley](#) | [BibTeX](#) |